



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Toward Standardized Monitoring of Patients With Chronic Diseases in Primary Care Using Electronic Medical Records: Systematic Review

Falck, Leandra ; Zoller, Marco ; Rosemann, Thomas ; Martínez-González, Nahara Anani ; Chmiel, Corinne

Abstract: BACKGROUND Long-term care for patients with chronic diseases poses a huge challenge in primary care. In particular, there is a deficit regarding monitoring and structured follow-up. Appropriate electronic medical records (EMRs) could help improving this but, so far, there are no evidence-based specifications concerning the indicators that should be monitored at regular intervals. OBJECTIVE The aim was to identify and collect a set of evidence-based indicators that could be used for monitoring chronic conditions at regular intervals in primary care using EMRs. METHODS We searched MEDLINE (Ovid), Embase (Elsevier), the Cochrane Library (Wiley), the reference lists of included studies and relevant reviews, and the content of clinical guidelines. We included primary studies and guidelines reporting about indicators that allow for the assessment of care and help monitor the status and process of disease for five chronic conditions, including type 2 diabetes mellitus, asthma, arterial hypertension, chronic heart failure, and osteoarthritis. RESULTS The use of the term "monitoring" in terms of disease management and long-term care for patients with chronic diseases is not widely used in the literature. Nevertheless, we identified a substantial number of disease-specific indicators that can be used for routine monitoring of chronic diseases in primary care by means of EMRs. CONCLUSIONS To our knowledge, this is the first systematic review summarizing the existing scientific evidence on the standardized long-term monitoring of chronic diseases using EMRs. In a second step, our extensive set of indicators will serve as a generic template for evaluating their usability by means of an adapted Delphi procedure. In a third step, the indicators will be summarized into a user-friendly EMR layout.

DOI: <https://doi.org/10.2196/10879>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-171182>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Falck, Leandra; Zoller, Marco; Rosemann, Thomas; Martínez-González, Nahara Anani; Chmiel, Corinne (2019). Toward Standardized Monitoring of Patients With Chronic Diseases in Primary Care Using Electronic Medical Records: Systematic Review. *JMIR Medical Informatics*, 7(2):e10879.

DOI: <https://doi.org/10.2196/10879>

Review

Toward Standardized Monitoring of Patients With Chronic Diseases in Primary Care Using Electronic Medical Records: Systematic Review

Leandra Falck, MD; Marco Zoller, MD; Thomas Rosemann, MD, PhD; Nahara Anani Martínez-González, MSc; Corinne Chmiel, MD, PD

Institute of Primary Care, University of Zurich and University Hospital of Zurich, Zurich, Switzerland

Corresponding Author:

Leandra Falck, MD

Institute of Primary Care

University of Zurich and University Hospital of Zurich

Pestalozzistrasse 24

Zurich,

Switzerland

Phone: 41 44 255 98 55

Email: Leandra.Falck@usz.ch

Abstract

Background: Long-term care for patients with chronic diseases poses a huge challenge in primary care. In particular, there is a deficit regarding monitoring and structured follow-up. Appropriate electronic medical records (EMRs) could help improving this but, so far, there are no evidence-based specifications concerning the indicators that should be monitored at regular intervals.

Objective: The aim was to identify and collect a set of evidence-based indicators that could be used for monitoring chronic conditions at regular intervals in primary care using EMRs.

Methods: We searched MEDLINE (Ovid), Embase (Elsevier), the Cochrane Library (Wiley), the reference lists of included studies and relevant reviews, and the content of clinical guidelines. We included primary studies and guidelines reporting about indicators that allow for the assessment of care and help monitor the status and process of disease for five chronic conditions, including type 2 diabetes mellitus, asthma, arterial hypertension, chronic heart failure, and osteoarthritis.

Results: The use of the term “monitoring” in terms of disease management and long-term care for patients with chronic diseases is not widely used in the literature. Nevertheless, we identified a substantial number of disease-specific indicators that can be used for routine monitoring of chronic diseases in primary care by means of EMRs.

Conclusions: To our knowledge, this is the first systematic review summarizing the existing scientific evidence on the standardized long-term monitoring of chronic diseases using EMRs. In a second step, our extensive set of indicators will serve as a generic template for evaluating their usability by means of an adapted Delphi procedure. In a third step, the indicators will be summarized into a user-friendly EMR layout.

(*JMIR Med Inform* 2019;7(2):e10879) doi:[10.2196/10879](https://doi.org/10.2196/10879)

KEYWORDS

monitoring of chronic diseases; indicators; primary care; systematic review; electronic medical record; diabetes mellitus type 2; arterial hypertension; asthma; osteoarthritis; chronic heart failure

Introduction

In 2016, the World Health Organization estimated that 71% of the overall deaths worldwide occurred due to noncommunicable diseases [1]. The majority of these diseases include cardiovascular diseases, chronic respiratory diseases, and diabetes. In particular, the prevalence of type 2 (non-insulin-dependent) diabetes mellitus, arterial hypertension,

asthma, chronic heart failure, and musculoskeletal diseases is increasing rapidly around the world leading to increased multimorbidity and polypharmacy, especially in the older population [1,2]. The burden of these diseases consequently imposes a significant threat to health, quality of life, and economic status in the affected population. Moreover, the regular monitoring of chronic diseases poses huge challenges and requires knowledge and communication skills, as well as the

capability of organization and coordination. The chronic care model (CCM) was originally introduced to graphically picture the concept of disease management [3]. The eHealth enhanced chronic care model was subsequently introduced as the means to improve the CCM in view of the progress and development of information and communication technology [4]. This model shows the existing variety of technically well-advanced applications as part of the monitoring process. Too many clinical offices in Switzerland lack basic electronic devices since many general practitioners still use paper-based patient records.

In 2012, 31 European countries were ranked based on the usage of electronic medical records (EMRs) in primary care [5]. In this global ranking of EMR usage, Switzerland ranked number 24. In a Swiss study, only up to 44.8% of the participating primary care physicians reported the usage of EMRs [6]. Therefore, it is currently almost impossible to exchange data with digital applications that are increasingly available and used by patients [6]. To efficiently monitor patients with chronic diseases, a well-structured and organized EMR system is crucial to ensure that all necessary information can be easily entered and retrieved, while no essential information is missed. Surprisingly, there are no evidence-based specifications concerning the indicators that should be monitored at regular intervals. On one hand, there are currently no international standards for the monitoring of patients with chronic diseases by means of EMR in primary care. On the other hand, there are deficits regarding the actual monitoring and structured follow-up. Therefore, we aimed to identify and collect a set of evidence-based indicators that could be used for monitoring patients with chronic conditions at regular intervals in primary care using EMRs.

Methods

Systematic Identification and Assessment of Supporting Evidence

We followed the principles of systematic reviews [7] and developed a protocol a priori to guide the identification and assessment of the monitoring indicators.

Inclusion Criteria

We included clinical guidelines and primary peer-reviewed studies of any design, carried-out mainly in primary care (ie, family health care) patients aged 18 years and older, who were diagnosed with type 2 (non-insulin-dependent) diabetes mellitus, arterial hypertension, asthma, chronic heart failure, or osteoarthritis. The first four diseases are among the most common noninfectious diseases worldwide. Osteoarthritis, in particular, generates a large part of indirect costs [2]. In order to be included, studies must have also reported on indicators that allow the assessment of care and help monitor the status and process of disease for these five chronic conditions. Therefore, we considered disease indicators that help reduce the risk of exacerbation, such as intermediate outcome indicators (eg, hemoglobin A_{1c} [HbA_{1c}] for diabetics or blood pressure measurements for hypertensive patients) and process indicators (eg, regular foot care or nutrition counselling). We included studies regardless of whether specific interventions were

evaluated. In addition, all studies and clinical guidelines should have been published in English or German.

Search Methods and Study Identification

We developed a comprehensive search strategy in collaboration with an expert librarian. The librarian conducted the search and produced a set of studies that matched the predefined search criteria. We identified studies published between 2000 and 2015 by applying this strategy in MEDLINE (Ovid), Embase (Elsevier), and the Cochrane Library (Wiley). No restrictions were made regarding the country of origin of the studies. The search strategy included a combination of the concepts and terminology, synonyms and related words for monitoring and for medical, health, electronic, patient, or file records. It also included primary, family, health care, or general practitioner, and the five chronic conditions (ie, type 2 [non-insulin-dependent] diabetes mellitus, arterial hypertension, asthma, chronic heart failure, and osteoarthritis). The focused search also included the terminology indicators, parameter, and management. An example of the full search strategy is available in [Multimedia Appendix 1](#).

We identified additional publications by manually searching the reference lists of included studies and relevant reviews. We also searched for monitoring indicators in the clinical guidelines in order to identify as many indicators as possible and to enable a holistic management of chronic diseases. Given that most guidelines are not indexed in the former medical literature databases, and to identify the clinical guidelines related to any of the five chronic diseases, we searched World Wide Web-based databases, including the National Guideline Clearinghouse for US guidelines [8] and the Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften eV (AWMF) [9] for German guidelines.

Study Selection and Assessment

For study selection, we created a system to prioritize the studies. One reviewer identified eligible studies by first screening the titles and abstracts of all records retrieved by the searches based on the inclusion criteria. All potentially eligible abstracts were rated manually from one to five stars according to their relevance for this review. The stars were assigned based on whether or not the key terms were mentioned (ie, “indicator,” “monitoring,” “assessment,” “management,” and/or “guideline”). The ranking was assigned as follows:

1. One star: Remote reference to the key terms; no indicators expected in full text.
2. Two stars: Little reference to the key terms; indicators in full text unlikely.
3. Three stars: Reference of at least one key term; indicators in full text possible.
4. Four stars: Reference of at least one key term; indicators in full text very possible.
5. Five stars: Reference of indicators, monitoring, or interval of measuring indicators.

The full text of all studies with an abstract that was rated with at least two stars was obtained, if available, and further evaluated based on the reporting of indicators. For studies where the full text was not available but were deemed important to inform our

monitoring tool, we used the data reported in the abstract. When it was necessary, the study team was consulted throughout the evaluation process to confirm the eligibility of indicators.

Data Extraction and Synthesis


For each included study, we extracted the bibliographic details (ie, author, year, and country of origin), all the monitoring indicators reported, the guideline on which the indicators were based, and the country of origin of the guidelines for each of the five chronic diseases. One reviewer extracted all data, and another reviewer verified the extracted data. We compiled a data profile for each study or guideline, and generated a set of

indicators using Microsoft Excel. We report a descriptive summary of the indicators for each of the chronic conditions.

Results

Our literature searches identified 795 original records (see Figure 1). After deduplication and perusal of titles and abstracts, we screened 621 records (range by disease: 33-180) and excluded 408 records that did not meet our inclusion criteria (eg, focused on specific therapy or medication or did not cover the topic). We examined in detail the full text, where available, of 213 publications (range by disease: 13 to 82).

Figure 1. Flowchart demonstrating the identification and selection of evidence. a: type 2 diabetes mellitus; b: asthma; c: arterial hypertension; d: heart failure; e: osteoarthritis; *: 5 of 87 publications (6%) reported indicators for more than one disease of interest.



We included 87 original publications, 5 (6%) in abstract form only, reporting indicators for diabetes mellitus [10-63], asthma [60,64-70], arterial hypertension [10,35,39,71-81], heart failure [33,82-92], and osteoarthritis [93-96]. [Multimedia Appendix 2](#) presents a list of all included studies that reported monitoring indicators for the five chronic conditions. A total of 5 publications (6%) reported indicators for more than one chronic disease [10,33,35,39,60]. The number of included publications by disease with at least one indicator ranged from 4 to 54. Most records (54/87, 62%) were published on type 2 diabetes mellitus, while osteoarthritis was the most underrepresented of the five diseases, with only 4 records (5%). A total of 74 of all 87

included studies (85%) contained process indicators, the most significant type of indicators. Concerning diabetes mellitus, a third of all publications (54/179, 30.2%) reported at least one indicator. For arterial hypertension and heart failure, only 8% (7/87) of all publications reported at least one indicator. Overall, most records used guidelines from the United States, followed by the United Kingdom. For diabetes mellitus, the American Diabetes Association and the National Institute for Health and Care Excellence were the most-used guidelines. The most frequently mentioned indicators for diabetes are presented in [Table 1](#). The indicators for the other four diseases are presented in [Multimedia Appendices 3-6](#).

Table 1. Diabetes mellitus indicators that are most frequently mentioned in guidelines and studies. The indicators are sorted first by guidelines and then by studies.

Indicators for diabetes mellitus	Number of guidelines where indicators are mentioned (guidelines)	Number of studies where indicators are mentioned
Fundoscopy examination	7 (a-g) ^a	20 [10,12,13,18,21,23,25,30-32,41,43-46,50-52,60,61]
Height, weight, and body mass index	7 (a-g)	33 [11-16,18,20,21,23-25,27-29,32,33,35,40,41,44,45,48,49,53-56,58-60,62,63]
Blood pressure measurement	7 (a-g)	45 [11,13,15-27,29-36,39-45,47-49,52,53,55,56,58-63]
10 g monofilament	7 (a-g)	N/A ^b
Hemoglobin A _{1c} (ie, glycated hemoglobin)	7 (a-g)	46 [10,12,13,15-23,26,28-37,39-45,47-54,56-63]
Foot inspection	7 (a-g)	17 [12,15,18,21,23,25,30-32,43-46,50-52,61]
Erectile dysfunction	7 (a-g)	N/A
Albuminuria	7 (a-g)	18 [12,13,18,22,23,25,31,32,35,41,43-46,51,55,61,62]
Lipid profile	7 (a-g)	8 [25,26,30,43,45,46,52,61]
Low-density lipoprotein	N/A	30 [11,12,15,18-20,22-24,29,31-37,41-44,47-49,52-54,63]
High-density lipoprotein	N/A	14 [11,20,23,28,29,33,37,39,49,51,53,54,62,63]
Triglyceride	N/A	15 [20,29,30,33,37,39,48,49,51,53-55,57,62,63]
Creatinine	7 (a-g)	18 [13,15,16,22,25-27,29,33,41,46,51,55,57-60,62]
Alcohol intake	7 (a-g)	2 [24,53]
Neuropathy and history of foot lesion	7 (a-g)	3 [18,20,55]
History of myocardial infarction (ie, cardiovascular disease)	6 (a-f)	2 [18,22]
Foot pulses	6 (a-f)	3 [18,32,60]
Smoking status	6 (a-f)	24 [11-15,18,20,22-24,26,28,29,31,35,41,44,48,50,53,58-61]
Orthostatic hypotension	5 (a, b, d, e, g)	N/A
Skin inspection	5 (a, b, d, f, g)	N/A
Vibration by 128 Hz tuning fork	5 (a-d, g)	1 [60]
Plasma glucositis	4 (b-d, g)	12 [11,21,24,33,39,45,51,54,55,57,59,63]
Onset of diabetes	3 (b, c, f)	9 [11,18,22,23,28,48,55,58,59]
Indicators appeared in fewer than five guidelines	225	N/A
Indicators appeared in fewer than 10 studies	N/A	76

^aThe letters a-g refer to the guidelines listed in [Multimedia Appendices 7-11](#).

^bN/A: not applicable.

In total, there were 249 indicators for type 2 diabetes mellitus, 183 for asthma, 335 for arterial hypertension, 231 for chronic heart failure, and 164 for osteoarthritis. The majority of indicators were identified by screening both peer-reviewed articles and clinical guidelines. A few extra indicators were reported only in peer-reviewed articles. That is, clinical guidelines on their own contributed to the great majority of all indicators identified. Surprisingly, only a few guidelines, such as the American guideline for asthma, included a section dedicated to monitoring or follow-up. Most of the guidelines that we screened did not specify the interval at which the indicators should be monitored. Also, in some guidelines, self-monitoring was a big topic for chronic heart disease (ie, weight control), asthma (ie, peak expiratory flow), and type 2 diabetes mellitus (ie, glucose monitoring).

Our systematic review also found that the term “monitoring,” in the sense of long-term patient care, was not widely used. Although publications reported the actual monitoring indicators, the process of monitoring for the different diseases, including, for example, the potential risks associated with overmonitoring, was only scarcely addressed. The publication by Glasziou was the only one giving a broader overview on the topic [97]. Only a handful of publications reported a complete set of indicators that can be used for monitoring, but these were either not specific for primary care or not eligible for implementation in EMRs [98-101].

Discussion

Principal Findings

To our knowledge, this study represents the first summary of the existing scientific evidence about the indicators that help standardize the monitoring of chronically ill patients in primary care by the use of EMRs. Long-term care of patients with chronic diseases is challenging and there are deficits regarding their monitoring and structured follow-up. Chronic care often involves collaboration between several people involved in the treatment process. That is only one reason for its complexity. Interpersonal differences in monitoring can decrease the quality of monitoring processes. Surprisingly, there are currently no gold standards or consensus regarding the systematic monitoring of patients with chronic diseases, in particular by means of EMRs. To efficiently monitor patients with chronic diseases, a well-structured and organized EMR system is crucial to ensure that all necessary information can be easily entered and retrieved and that no essential information is missed. Our study is, thus, the first initiative toward the urgent need of standardization for monitoring patients with chronic diseases in primary care.

Our systematic literature review showed that the term “monitoring” in terms of disease management and long-term patient care is not widely used. There is a plethora of literature about quality indicators that might have the potential to improve the outcome of a disease. The Quality and Outcomes Framework (QOF) in the United Kingdom, for example, assesses indicators for such purposes [102]. Beyond identifying indicators that can be easily assessed, such as the indicators used by the QOF, our goal was to summarize the existing literature on all the indicators available for long-term monitoring.

So far, only a few authors have focused on the topic of the monitoring of chronic diseases. According to Glasziou, the process of monitoring aims to establish the response to treatment and to detect both adverse effects and the need to adjust treatment [97]. The process of monitoring can be divided into different phases (ie, pretreatment, during treatment, and after treatment). Each phase requires measurements at different intervals.

When analyzing different diseases, monitoring is probably most widely mentioned in blood pressure management. There are various publications reporting on the optimal way and interval of measuring blood pressure [76,103,104]. However, literature beyond the indicator of blood pressure measurement remains scarce. Regarding diabetes mellitus, there is an extended monitoring tool that was designed as a disease management tool for practice nurses [101]. The tool’s design is based on a traffic light scheme to detect any deficit and need for action. In addition, a detailed guideline on how to monitor the diabetic foot is provided by the International Working Group on the Diabetic Foot [105]. As for bronchial asthma, two study groups have addressed the optimal way and potential problems of finding and evaluating indicators to monitor patients with asthma, including an overview of the most important indicators [98,100]. Similarly, Grypdonck presents a small set of indicators for monitoring patients with osteoarthritis of the knee [93]. Self-monitoring seems to be an important topic concerning osteoarthritis and asthma. An English study conducted by interviewing general practitioners about osteoarthritis showed that the majority of respondents thought monitoring of osteoarthritis is important, even though almost half did not monitor patients at all. Interestingly, more than half of the respondents felt that patients should do self-monitoring [106]. Patient involvement is crucial for monitoring. Particularly, in high-frequency monitoring situations such as chronic heart failure, telecardiological service, including transtelephonic monitoring, reduces the length of hospitalization and improves quality of life [91]. Surprisingly, publications concerning monitoring of chronic heart failure seem to be scarce [90]. The underrepresentation of osteoarthritis and chronic heart failure is also reflected in the number of indicators detected in the primary literature, compared to a large number of records reporting on indicators for type 2 diabetes mellitus. Another topic repeatedly found in the results was the involvement of a clinical practice nurse in monitoring [101,107-109]. The clinical nurse can, for example, fill out a monitoring questionnaire in face-to-face sessions with the patient, on the phone, or even electronically. This could counteract the problem of workload and time constraints as a frequent response to why monitoring is not conducted [106].

Strengths and Limitations

To our knowledge, this study represents the first scientifically founded recommendation for the standardized long-term monitoring of chronically ill patients in primary care. Usually, systematic reviews only concentrate on primary literature and do not include guidelines in their search strategy, since most guidelines are not indexed in databases. In our study, we explicitly searched for guideline programs such as the National Guideline Clearinghouse for American guidelines and the

AWMF for German guidelines. We added a substantial number of manual searches within reference lists and search engines in order to gain a maximal insight of the existing literature. This strategy was worth the extra effort, considering that most relevant indicators were found in guidelines and not in the primary literature. Possible confounders are that publications and guidelines reported in languages other than German and English were excluded.

Outlook

In a second step, our extensive set of indicators obtained from this work will serve as a generic template for a monitoring tool.

By means of an adapted Delphi procedure, the indicators will be further evaluated in terms of their usability. In a third step, the indicators will be summarized into a user-friendly EMR layout.

Conclusion

This is the first study that systematically summarizes the existing scientific evidence about the standardized long-term monitoring of chronic diseases by means of EMRs. It aims to help improve care for patients with chronic diseases in primary care.

Acknowledgments

We are grateful to Martina Gosteli for her assistance with the search strategy. This project was funded by the Institut für Praxisinformatik, mandated by the Swiss Medical Association FMH.

Authors' Contributions

LF was involved in designing the search strategy, performing the systematic screening and review of the literature, and writing of the manuscript. MZ was involved in study design and revised the manuscript. NAMG contributed to the study design and search strategy; NAMG also revised and improved the manuscript. TR supervised the development and methodology of the study and helped improve the final version of the manuscript. CC was involved in the study design, study selection, and prioritization; CC also verified the extracted data, and supervised and revised the manuscript. All authors read and approved the final manuscript.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Search strategy for MEDLINE (Ovid), Embase (Elsevier), and the Cochrane Library (Wiley).

[\[DOCX File, 14KB - medinform_v7i2e10879_app1.docx\]](#)

Multimedia Appendix 2

List of all included studies, including monitoring indicators for the five chronic conditions.

[\[DOCX File, 27KB - medinform_v7i2e10879_app2.docx\]](#)

Multimedia Appendix 3

Asthma indicators mentioned in guidelines and studies.

[\[DOCX File, 15KB - medinform_v7i2e10879_app3.docx\]](#)

Multimedia Appendix 4

Arterial hypertension indicators mentioned in guidelines and studies.

[\[DOCX File, 16KB - medinform_v7i2e10879_app4.docx\]](#)

Multimedia Appendix 5

Chronic heart failure indicators most frequently mentioned in guidelines and studies.

[\[DOCX File, 14KB - medinform_v7i2e10879_app5.docx\]](#)

Multimedia Appendix 6

Osteoarthritis indicators most frequently mentioned in guidelines and studies.

[\[DOCX File, 14KB - medinform_v7i2e10879_app6.docx\]](#)

Multimedia Appendix 7

Guidelines screened for indicators for type 2 diabetes mellitus.

[[DOCX File, 14KB](#) - [medinform_v7i2e10879_app7.docx](#)]

Multimedia Appendix 8

Guidelines screened for indicators for asthma.

[[DOCX File, 13KB](#) - [medinform_v7i2e10879_app8.docx](#)]

Multimedia Appendix 9

Guidelines screened for indicators for arterial hypertension.

[[DOCX File, 14KB](#) - [medinform_v7i2e10879_app9.docx](#)]

Multimedia Appendix 10

Guidelines screened for indicators for chronic heart failure.

[[DOCX File, 14KB](#) - [medinform_v7i2e10879_app10.docx](#)]

Multimedia Appendix 11

Guidelines screened for indicators for osteoarthritis.

[[DOCX File, 13KB](#) - [medinform_v7i2e10879_app11.docx](#)]

References

1. World Health Statistics 2018: Monitoring Health for the Sustainable Development Goals (SDGs). Geneva, Switzerland: World Health Organization; 2018. URL: https://www.who.int/gho/publications/world_health_statistics/2016/Annex_B/en/ [accessed 2019-05-14] [WebCite Cache ID 78Mf6qAst]
2. Bachmann N, Burla L, Kohler D, Schweizerisches Gesundheitsobservatorium. Gesundheit in der Schweiz: Fokus Chronische Erkrankungen. Nationaler Gesundheitsbericht 2015. Bern, Switzerland: Hogrefe Verlag; 2015. URL: https://www.obsan.admin.ch/sites/default/files/publications/2015/gesundheitsbericht_2015_d.pdf [accessed 2019-05-13] [WebCite Cache ID 78LZfvcWG]
3. Wagner EH. Chronic disease management: What will it take to improve care for chronic illness? Eff Clin Pract 1998;1(1):2-4. [Medline: 10345255]
4. Gee PM, Greenwood DA, Paterniti DA, Ward D, Miller LM. The eHealth Enhanced Chronic Care Model: A theory derivation approach. J Med Internet Res 2015 Apr 01;17(4):e86 [FREE Full text] [doi: 10.2196/jmir.4067] [Medline: 25842005]
5. De Rosi S, Seghieri C. Basic ICT adoption and use by general practitioners: An analysis of primary care systems in 31 European countries. BMC Med Inform Decis Mak 2015 Aug 22;15:70 [FREE Full text] [doi: 10.1186/s12911-015-0185-z] [Medline: 26296994]
6. Djalali S, Ursprung N, Rosemann T, Senn O, Tandjung R. Undirected health IT implementation in ambulatory care favors paper-based workarounds and limits health data exchange. Int J Med Inform 2015 Nov;84(11):920-932. [doi: 10.1016/j.ijmedinf.2015.08.001] [Medline: 26296433]
7. Egger M, Smith GD. Principles of and procedures for systematic reviews. In: Egger M, Smith GD, Altman DG, editors. Systematic Reviews in Health Care: Meta-Analysis in Context. 2nd edition. London, UK: BMJ Publishing Group; 2008:23-42.
8. Agency for Healthcare Research and Quality. 2015. National Guideline Clearinghouse URL: <https://www.guideline.gov/> [WebCite Cache ID 6yobWTV23]
9. AWMF Das Portal der wissenschaftlichen Medizin. Leitliniensuche URL: <http://www.awmf.org> [WebCite Cache ID 6yodPI7r6]
10. Suija K, Kivisto K, Sarria-Santamera A, Kokko S, Liseckiene I, Bredehorst M, et al. Challenges of audit of care on clinical quality indicators for hypertension and type 2 diabetes across four European countries. Fam Pract 2015 Feb;32(1):69-74. [doi: 10.1093/fampra/cmu078] [Medline: 25411423]
11. Shah AD, Langenberg C, Rapsomaniki E, Denaxas S, Pujades-Rodriguez M, Gale CP, et al. Type 2 diabetes and incidence of cardiovascular diseases: A cohort study in 1·9 million people. Lancet Diabetes Endocrinol 2015 Feb;3(2):105-113 [FREE Full text] [doi: 10.1016/S2213-8587(14)70219-0] [Medline: 25466521]
12. Devkota BP, Anstas M, Scherrer JF, Salas J, Budhathoki C. Internal medicine resident training and provision of diabetes quality of care indicators. Can J Diabetes 2015 Apr;39(2):133-137. [doi: 10.1016/j.cjcd.2014.10.001] [Medline: 25499028]

13. Barkhuysen P, de Grauw W, Akkermans R, Donkers J, Schers H, Biermans M. Is the quality of data in an electronic medical record sufficient for assessing the quality of primary care? *J Am Med Inform Assoc* 2014;21(4):692-698 [[FREE Full text](#)] [doi: [10.1136/amiajnl-2012-001479](#)] [Medline: [24145818](#)]
14. Szczech LA, Stewart RC, Su H, DeLoskey RJ, Astor BC, Fox CH, et al. Primary care detection of chronic kidney disease in adults with type-2 diabetes: The ADD-CKD Study (awareness, detection and drug therapy in type 2 diabetes and chronic kidney disease). *PLoS One* 2014;9(11):e110535 [[FREE Full text](#)] [doi: [10.1371/journal.pone.0110535](#)] [Medline: [25427285](#)]
15. van Melle MA, Lamkaddem M, Stuiver MM, Gerritsen AA, Devillé WL, Essink-Bot ML. Quality of primary care for resettled refugees in the Netherlands with chronic mental and physical health problems: A cross-sectional analysis of medical records and interview data. *BMC Fam Pract* 2014 Sep 23;15:160 [[FREE Full text](#)] [doi: [10.1186/1471-2296-15-160](#)] [Medline: [25249345](#)]
16. Djalali S, Frei A, Tandjung R, Baltensperger A, Rosemann T. Swiss quality and outcomes framework: Quality indicators for diabetes management in Swiss primary care based on electronic medical records. *Gerontology* 2014;60(3):263-273. [doi: [10.1159/000357370](#)] [Medline: [24603324](#)]
17. Goff SL, Murphy L, Knee AB, Guhn-Knight H, Guhn A, Lindenauer PK. Effects of an enhanced primary care program on diabetes outcomes. *Am J Manag Care* 2017 Mar 01;23(3):e75-e81 [[FREE Full text](#)] [Medline: [28385028](#)]
18. Vidal-Pardo JI, Pérez-Castro TR, López-Álvarez XL, Santiago-Pérez MI, García-Soidán FJ, Muñoz J. Effect of an educational intervention in primary care physicians on the compliance of indicators of good clinical practice in the treatment of type 2 diabetes mellitus [OBTEDIGA project]. *Int J Clin Pract* 2013 Aug;67(8):750-758. [doi: [10.1111/ijcp.12145](#)] [Medline: [23668834](#)]
19. Sidorenkov G, Voorham J, de Zeeuw D, Haaïjer-Ruskamp FM, Denig P. Treatment quality indicators predict short-term outcomes in patients with diabetes: A prospective cohort study using the GIANIT database. *BMJ Qual Saf* 2013 Apr;22(4):339-347. [doi: [10.1136/bmjqs-2012-001203](#)] [Medline: [23386729](#)]
20. Winkley K, Thomas SM, Sivaprasad S, Chamley M, Stahl D, Ismail K, et al. The clinical characteristics at diagnosis of type 2 diabetes in a multi-ethnic population: The South London Diabetes cohort (SOUL-D). *Diabetologia* 2013 Jun;56(6):1272-1281. [doi: [10.1007/s00125-013-2873-5](#)] [Medline: [23494447](#)]
21. Gavran L, Zildzic M, Batic-Mujanovic O, Alic A, Gledio I, Prasko S. Auditing of medical chart among type 2 diabetic patient done by primary care physicians. *Med Arch* 2012;66(6):388-390. [Medline: [23409518](#)]
22. Knudsen ST, Mosbech TH, Hansen B, König E, Johnsen PC, Kamper A. Screening for microalbuminuria in patients with type 2 diabetes is incomplete in general practice. *Dan Med J* 2012 Sep;59(9):A4502. [Medline: [22951198](#)]
23. Mata-Cases M, Roura-Olmeda P, Berengué-Iglesias M, Birulés-Pons M, Mundet-Tuduri X, Franch-Nadal J, Diabetes Study Group in Primary Health Care (GEDAPS: Grup d'Estudi de la Diabetis a l'Atenció Primària de Salut, Catalanian Society of Family and Community Medicine). Fifteen years of continuous improvement of quality care of type 2 diabetes mellitus in primary care in Catalonia, Spain. *Int J Clin Pract* 2012 Mar;66(3):289-298 [[FREE Full text](#)] [doi: [10.1111/j.1742-1241.2011.02872.x](#)] [Medline: [22340449](#)]
24. Nouwens E, van Lieshout J, Wensing M. Comorbidity complicates cardiovascular treatment: Is diabetes the exception? *Neth J Med* 2012 Sep;70(7):298-305 [[FREE Full text](#)] [Medline: [22961822](#)]
25. Satman I, Imamoglu S, Yilmaz C, ADMIRE Study Group. A patient-based study on the adherence of physicians to guidelines for the management of type 2 diabetes in Turkey. *Diabetes Res Clin Pract* 2012 Oct;98(1):75-82. [doi: [10.1016/j.diabres.2012.05.003](#)] [Medline: [22652276](#)]
26. Marley JV, Nelson C, O'Donnell V, Atkinson D. Quality indicators of diabetes care: An example of remote-area Aboriginal primary health care over 10 years. *Med J Aust* 2012 Oct 01;197(7):404-408. [Medline: [23025738](#)]
27. Patapas JM, Blanchard AC, Iqbal S, Vasilevsky M, Dannenbaum D. Management of aboriginal and nonaboriginal people with chronic kidney disease in Quebec: Quality-of-care indicators. *Can Fam Physician* 2012 Feb;58(2):e107-e111 [[FREE Full text](#)] [Medline: [22439172](#)]
28. Staff M. Can data extraction from general practitioners' electronic records be used to predict clinical outcomes for patients with type 2 diabetes? *Inform Prim Care* 2012;20(2):95-102 [[FREE Full text](#)] [Medline: [23710774](#)]
29. Hill F, Bradley C. A computer based, automated analysis of process and outcomes of diabetic care in 23 GP practices. *Ir Med J* 2012 Feb;105(2):45-47. [Medline: [22455238](#)]
30. Alfadda AA, Bin-Abdulrahman KA, Saad HA, Mendoza CD, Angkaya-Bagayawa FF, Yale JF. Effect of an intervention to improve the management of patients with diabetes in primary care practice. *Saudi Med J* 2011 Jan;32(1):36-40. [Medline: [21212914](#)]
31. Dickerson LM, Ables AZ, Everett CJ, Mainous AG, McCutcheon AM, Bazaldua OV, National Interdisciplinary Primary Care Practice-Based Research Network. Measuring diabetes care in the national interdisciplinary primary care practice-based research network (NIPC-PBRN). *Pharmacotherapy* 2011 Jan;31(1):23-30. [doi: [10.1592/phco.31.1.23](#)] [Medline: [21182355](#)]
32. Vidal Pardo JI, Pérez Castro TR, López Álvarez XL, García Soidán FJ, Santiago Pérez MI, Muñoz J. Quality of care of patients with type-2 diabetes in Galicia (NW Spain) [OBTEDIGA project]. *Int J Clin Pract* 2011 Oct;65(10):1067-1075. [doi: [10.1111/j.1742-1241.2011.02739.x](#)] [Medline: [21801286](#)]

33. Weenink J, van Lieshout J, Jung HP, Wensing M. Patient care teams in treatment of diabetes and chronic heart failure in primary care: An observational networks study. *Implement Sci* 2011 Jul 03;6:66 [[FREE Full text](#)] [doi: [10.1186/1748-5908-6-66](https://doi.org/10.1186/1748-5908-6-66)] [Medline: [21722399](#)]
34. Gladstone J, Howard M. Effect of advanced access scheduling on chronic health care in a Canadian practice. *Can Fam Physician* 2011 Jan;57(1):e21-e25 [[FREE Full text](#)] [Medline: [21252121](#)]
35. Holbrook A, Pullenayegum E, Thabane L, Troyan S, Foster G, Keshavjee K, et al. Shared electronic vascular risk decision support in primary care: Computerization of Medical Practices for the Enhancement of Therapeutic Effectiveness (COMPETE III) randomized trial. *Arch Intern Med* 2011 Oct 24;171(19):1736-1744. [doi: [10.1001/archinternmed.2011.471](https://doi.org/10.1001/archinternmed.2011.471)] [Medline: [22025430](#)]
36. O'Connor PJ, Sperl-Hillen JM, Rush WA, Johnson PE, Amundson GH, Asche SE, et al. Impact of electronic health record clinical decision support on diabetes care: A randomized trial. *Ann Fam Med* 2011;9(1):12-21 [[FREE Full text](#)] [doi: [10.1370/afm.1196](https://doi.org/10.1370/afm.1196)] [Medline: [21242556](#)]
37. Sundquist K, Chaikiat A, León VR, Johansson S, Sundquist J. Country of birth, socioeconomic factors, and risk factor control in patients with type 2 diabetes: A Swedish study from 25 primary health-care centres. *Diabetes Metab Res Rev* 2011 Mar;27(3):244-254. [doi: [10.1002/dmrr.1161](https://doi.org/10.1002/dmrr.1161)] [Medline: [21309045](#)]
38. Reddy P, Philpot B, Ford D, Dunbar JA. Identification of depression in diabetes: The efficacy of PHQ-9 and HADS-D. *Br J Gen Pract* 2010 Jun;60(575):e239-e245 [[FREE Full text](#)] [doi: [10.3399/bjgp10X502128](https://doi.org/10.3399/bjgp10X502128)] [Medline: [20529487](#)]
39. Samoutis GA, Soteriades ES, Stoffers HE, Philalithis A, Delicha EM, Lionis C. A pilot quality improvement intervention in patients with diabetes and hypertension in primary care settings of Cyprus. *Fam Pract* 2010 Jun;27(3):263-270. [doi: [10.1093/fampra/cmq009](https://doi.org/10.1093/fampra/cmq009)] [Medline: [20348163](#)]
40. Shah BR, Bhattacharyya O, Yu C, Mamdani M, Parsons JA, Straus SE, et al. Evaluation of a toolkit to improve cardiovascular disease screening and treatment for people with type 2 diabetes: Protocol for a cluster-randomized pragmatic trial. *Trials* 2010 Apr 23;11:44 [[FREE Full text](#)] [doi: [10.1186/1745-6215-11-44](https://doi.org/10.1186/1745-6215-11-44)] [Medline: [20416080](#)]
41. Petrazzuoli F, Soler JK, Buono N, Dobbs F. Quality of care for hypertensive patients with type 2 diabetes in a rural area of Southern Italy: Is the recording of patient data and the achievement of quality indicators targets satisfactory? *Rural Remote Health* 2010;10(3):1258 [[FREE Full text](#)] [Medline: [20843159](#)]
42. Sperl-Hillen JM, O'Connor PJ, Rush WA, Johnson PE, Gilmer T, Biltz G, et al. Simulated physician learning program improves glucose control in adults with diabetes. *Diabetes Care* 2010 Aug;33(8):1727-1733 [[FREE Full text](#)] [doi: [10.2337/dc10-0439](https://doi.org/10.2337/dc10-0439)] [Medline: [20668151](#)]
43. Pedersen ML. Management of type 2 diabetes mellitus in Greenland, 2008: Examining the quality and organization of diabetes care. *Int J Circumpolar Health* 2009 Apr;68(2):123-132. [Medline: [19517872](#)]
44. Holbrook A, Thabane L, Keshavjee K, Dolovich L, Bernstein B, Chan D, COMPETE II Investigators. Individualized electronic decision support and reminders to improve diabetes care in the community: COMPETE II randomized trial. *CMAJ* 2009 Jul 07;181(1-2):37-44 [[FREE Full text](#)] [doi: [10.1503/cmaj.081272](https://doi.org/10.1503/cmaj.081272)] [Medline: [19581618](#)]
45. Moharram MM, Farahat FM. Quality improvement of diabetes care using flow sheets in family health practice. *Saudi Med J* 2008 Jan;29(1):98-101. [Medline: [18176682](#)]
46. Novo A, Jokić I. Medical audit of diabetes mellitus in primary care setting in Bosnia and Herzegovina. *Croat Med J* 2008 Dec;49(6):757-762 [[FREE Full text](#)] [Medline: [19090600](#)]
47. Samuels TA, Bolen S, Yeh HC, Abuid M, Marinopoulos SS, Weiner JP, et al. Missed opportunities in diabetes management: A longitudinal assessment of factors associated with sub-optimal quality. *J Gen Intern Med* 2008 Nov;23(11):1770-1777 [[FREE Full text](#)] [doi: [10.1007/s11606-008-0757-z](https://doi.org/10.1007/s11606-008-0757-z)] [Medline: [18787908](#)]
48. Smith SA, Shah ND, Bryant SC, Christianson TJ, Bjornsen SS, Giesler PD, Evidens Research Group. Chronic care model and shared care in diabetes: Randomized trial of an electronic decision support system. *Mayo Clin Proc* 2008 Jul;83(7):747-757. [doi: [10.4065/83.7.747](https://doi.org/10.4065/83.7.747)] [Medline: [18613991](#)]
49. Voorham J, Denig P, Wolffenbuttel BH, Haaijer-Ruskamp FM. Cross-sectional versus sequential quality indicators of risk factor management in patients with type 2 diabetes. *Med Care* 2008 Feb;46(2):133-141. [doi: [10.1097/MLR.0b013e31815b9da0](https://doi.org/10.1097/MLR.0b013e31815b9da0)] [Medline: [18219241](#)]
50. Wens J, Dirven K, Mathieu C, Paulus D, Van Royen P, Belgian Diabetes Project Group. Quality indicators for type-2 diabetes care in practice guidelines: An example from six European countries. *Prim Care Diabetes* 2007 Feb;1(1):17-23. [doi: [10.1016/j.pcd.2006.07.001](https://doi.org/10.1016/j.pcd.2006.07.001)] [Medline: [18632015](#)]
51. Nitiyanant W, Chetthakul T, Sang-A-kad P, Therakiatkumjorn C, Kunsuikmengrai K, Yeo JP. A survey study on diabetes management and complication status in primary care setting in Thailand. *J Med Assoc Thai* 2007 Jan;90(1):65-71. [Medline: [17621735](#)]
52. Herrin J, Nicewander DA, Hollander PA, Couch CE, Winter FD, Haydar ZR, et al. Effectiveness of diabetes resource nurse case management and physician profiling in a fee-for-service setting: A cluster randomized trial. *Proc (Bayl Univ Med Cent)* 2006 Apr;19(2):95-102 [[FREE Full text](#)] [Medline: [16609732](#)]
53. Wan Q, Harris MF, Jayasinghe UW, Flack J, Georgiou A, Penn DL, et al. Quality of diabetes care and coronary heart disease absolute risk in patients with type 2 diabetes mellitus in Australian general practice. *Qual Saf Health Care* 2006 Apr;15(2):131-135 [[FREE Full text](#)] [doi: [10.1136/qshc.2005.014845](https://doi.org/10.1136/qshc.2005.014845)] [Medline: [16585115](#)]

54. Al Khaja KA, Sequeira RP, Damanhori AH. Comparison of the quality of diabetes care in primary care diabetic clinics and general practice clinics. *Diabetes Res Clin Pract* 2005 Nov;70(2):174-182. [doi: [10.1016/j.diabres.2005.03.029](https://doi.org/10.1016/j.diabres.2005.03.029)] [Medline: [15890429](#)]
55. Cueto-Manzano AM, Cortes-Sanabria L, Martinez-Ramirez HR, Rojas-Campos E, Barragan G, Alfaro G, et al. Detection of early nephropathy in Mexican patients with type 2 diabetes mellitus. *Kidney Int Suppl* 2005 Aug(97):S40-S45 [FREE Full text] [doi: [10.1111/j.1523-1755.2005.09707.x](https://doi.org/10.1111/j.1523-1755.2005.09707.x)] [Medline: [16014099](#)]
56. Lusignan S, Sismanidis C, Carey IM, DeWilde S, Richards N, Cook DG. Trends in the prevalence and management of diagnosed type 2 diabetes 1994-2001 in England and Wales. *BMC Fam Pract* 2005 Mar 22;6(1):13 [FREE Full text] [doi: [10.1186/1471-2296-6-13](https://doi.org/10.1186/1471-2296-6-13)] [Medline: [15784133](#)]
57. Sequeira RP, Al Khaja KA, Damanhori AH. Evaluating the treatment of hypertension in diabetes mellitus: A need for better control? *J Eval Clin Pract* 2004 Feb;10(1):107-116. [Medline: [14731158](#)]
58. Wermeille J, Bennie M, Brown I, McKnight J. Pharmaceutical care model for patients with type 2 diabetes: Integration of the community pharmacist into the diabetes team--A pilot study. *Pharm World Sci* 2004 Feb;26(1):18-25. [Medline: [15018255](#)]
59. Goudswaard AN, Lam K, Stolk RP, Rutten GE. Quality of recording of data from patients with type 2 diabetes is not a valid indicator of quality of care: A cross-sectional study. *Fam Pract* 2003 Apr;20(2):173-177. [Medline: [12651792](#)]
60. Campbell SM, Hann M, Hacker J, Durie A, Thapar A, Roland MO. Quality assessment for three common conditions in primary care: Validity and reliability of review criteria developed by expert panels for angina, asthma and type 2 diabetes. *Qual Saf Health Care* 2002 Jun;11(2):125-130 [FREE Full text] [doi: [10.1136/qhc.11.2.125](https://doi.org/10.1136/qhc.11.2.125)] [Medline: [12448803](#)]
61. Parchman ML, Burge SK, Residency Research Network of South Texas Investigators. Continuity and quality of care in type 2 diabetes: A Residency Research Network of South Texas study. *J Fam Pract* 2002 Jul;51(7):619-624. [Medline: [12160500](#)]
62. Renders CM, Valk GD, Franse LV, Schellevis FG, van Eijk JT, van der Wal G. Long-term effectiveness of a quality improvement program for patients with type 2 diabetes in general practice. *Diabetes Care* 2001 Aug;24(8):1365-1370. [Medline: [11473071](#)]
63. Linmans JJ, Spigt MG, Deneer L, Lucas AE, de Bakker M, Gidding LG, et al. Effect of lifestyle intervention for people with diabetes or prediabetes in real-world primary care: Propensity score analysis. *BMC Fam Pract* 2011 Sep 13;12:95 [FREE Full text] [doi: [10.1186/1471-2296-12-95](https://doi.org/10.1186/1471-2296-12-95)] [Medline: [21914190](#)]
64. Minard JP, Dostaler SM, Taite AK, Olajos-Clow JG, Sands TW, Licskai CJ, et al. Development and implementation of an electronic asthma record for primary care: Integrating guidelines into practice. *J Asthma* 2014 Feb;51(1):58-68. [doi: [10.3109/02770903.2013.845206](https://doi.org/10.3109/02770903.2013.845206)] [Medline: [24083321](#)]
65. Lim KG, Rank MA, Cabanela RL, Furst JW, Rohrer JE, Liesinger J, et al. The asthma ePrompt: A novel electronic solution for chronic disease management. *J Asthma* 2012 Mar;49(2):213-218. [doi: [10.3109/02770903.2012.654419](https://doi.org/10.3109/02770903.2012.654419)] [Medline: [22304226](#)]
66. Loughheed MD, Minard J, Dworkin S, Juurlink M, Temple WJ, To T, et al. Pan-Canadian REspiratory STandards INitiative for Electronic Health Records (PRESTINE): 2011 national forum proceedings. *Can Respir J* 2012;19(2):117-126 [FREE Full text] [doi: [10.1155/2012/870357](https://doi.org/10.1155/2012/870357)] [Medline: [22536581](#)]
67. Oei SM, Thien FC, Schattner RL, Sulaiman ND, Birch K, Simpson P, et al. Effect of spirometry and medical review on asthma control in patients in general practice: A randomized controlled trial. *Respirology* 2011 Jul;16(5):803-810. [doi: [10.1111/j.1440-1843.2011.01969.x](https://doi.org/10.1111/j.1440-1843.2011.01969.x)] [Medline: [21401801](#)]
68. Nokela M, Arnlind MH, Ehlers P, Krakau I, Forslund L, Jonsson EW. The influence of structured information and monitoring on the outcome of asthma treatment in primary care: A cluster randomized study. *Respiration* 2010;79(5):388-394 [FREE Full text] [doi: [10.1159/000235548](https://doi.org/10.1159/000235548)] [Medline: [19672055](#)]
69. Yawn BP, Bertram S, Wollan P. Introduction of Asthma APGAR tools improve asthma management in primary care practices. *J Asthma Allergy* 2008 Aug 31;1:1-10 [FREE Full text] [Medline: [21436980](#)]
70. Baddar S, Worthing EA, Al-Rawas OA, Osman Y, Al-Riyami BM. Compliance of physicians with documentation of an asthma management protocol. *Respir Care* 2006 Dec;51(12):1432-1440 [FREE Full text] [Medline: [17134524](#)]
71. Hasselström J, Zarrinkoub R, Holmquist C, Hjerpe P, Ljungman C, Qvarnström M, et al. The Swedish Primary Care Cardiovascular Database (SPCCD): 74 751 hypertensive primary care patients. *Blood Press* 2014 Apr;23(2):116-125. [doi: [10.3109/08037051.2013.814829](https://doi.org/10.3109/08037051.2013.814829)] [Medline: [23914944](#)]
72. Tong SF, Khoo EM, Nordin S, Teng C, Lee VK, Zailinawati AH, et al. Process of care and prescribing practices for hypertension in public and private primary care clinics in Malaysia. *Asia Pac J Public Health* 2012 Sep;24(5):764-775. [doi: [10.1177/1010539511402190](https://doi.org/10.1177/1010539511402190)] [Medline: [21659332](#)]
73. Pavlik VN, Greisinger AJ, Pool J, Haidet P, Hyman DJ. Does reducing physician uncertainty improve hypertension control? Rationale and methods. *Circ Cardiovasc Qual Outcomes* 2009 May;2(3):257-263 [FREE Full text] [doi: [10.1161/CIRCOUTCOMES.109.849984](https://doi.org/10.1161/CIRCOUTCOMES.109.849984)] [Medline: [20031846](#)]
74. Chan PF, Chao DVK. The Hong Kong Practitioner. 2006 Dec. An audit on management of hypertension in a hospital authority general outpatient clinic URL: http://www.hkcfp.org.hk/Upload/HK_Practitioner/2006/hkp2006vol28dec/original_article.html [accessed 2019-05-13] [WebCite Cache ID 78LcpOKYm]

75. Asnani M, Brown P, O'Connor D, Lewis T, Win S, Reid M. A clinical audit of the quality of care of hypertension in general practice. *West Indian Med J* 2005 Jun;54(3):176-180. [Medline: [16209222](#)]
76. Rabinowitz I, Tamir A. The SaM (Screening and Monitoring) approach to cardiovascular risk-reduction in primary care: Cyclic monitoring and individual treatment of patients at cardiovascular risk using the electronic medical record. *Eur J Cardiovasc Prev Rehabil* 2005 Feb;12(1):56-62. [Medline: [15703507](#)]
77. Mitchell E, Sullivan F, Grimshaw JM, Donnan PT, Watt G. Improving management of hypertension in general practice: A randomised controlled trial of feedback derived from electronic patient data. *Br J Gen Pract* 2005 Feb;55(511):94-101 [[FREE Full text](#)] [Medline: [15720929](#)]
78. Alli C, Mariotti G, Avanzini F, Colombo F, Barlera S, Tognoni G, Studio sulla Pressione Arteriosa nell'Anziano (SPAA). Long-term prognostic impact of repeated measurements over 1 year of pulse pressure and systolic blood pressure in the elderly. *J Hum Hypertens* 2005 May;19(5):355-363. [doi: [10.1038/sj.jhh.1001827](#)] [Medline: [15772693](#)]
79. Tierney WM, Brunt M, Kesterson J, Zhou X, L'Italien G, Lapuerta P. Quantifying risk of adverse clinical events with one set of vital signs among primary care patients with hypertension. *Ann Fam Med* 2004;2(3):209-217 [[FREE Full text](#)] [Medline: [15209196](#)]
80. Lackland DT, Lin Y, Tilley BC, Egan BM. An assessment of racial differences in clinical practices for hypertension at primary care sites for medically underserved patients. *J Clin Hypertens (Greenwich)* 2004 Jan;6(1):26-31; quiz 32 [[FREE Full text](#)] [Medline: [14724421](#)]
81. Frijling BD, Lobo CM, Hulscher ME, Akkermans RP, van Drenth BB, Prins A, et al. Intensive support to improve clinical decision making in cardiovascular care: A randomised controlled trial in general practice. *Qual Saf Health Care* 2003 Jun;12(3):181-187 [[FREE Full text](#)] [doi: [10.1136/qhc.12.3.181](#)] [Medline: [12792007](#)]
82. Amarasingham R, Patel PC, Toto K, Nelson LL, Swanson TS, Moore BJ, et al. Allocating scarce resources in real-time to reduce heart failure readmissions: A prospective, controlled study. *BMJ Qual Saf* 2013 Dec;22(12):998-1005 [[FREE Full text](#)] [doi: [10.1136/bmjqs-2013-001901](#)] [Medline: [23904506](#)]
83. Logeart D, Isnard R, Resche-Rigon M, Seronde M, de Groote P, Jondeau G, Heart Failure of the French Society of Cardiology. Current aspects of the spectrum of acute heart failure syndromes in a real-life setting: The OFICA study. *Eur J Heart Fail* 2013 Apr;15(4):465-476 [[FREE Full text](#)] [doi: [10.1093/eurjhf/hfs189](#)] [Medline: [23186936](#)]
84. Lind L, Klompstra L, Jaarsma T, Strömberg A. Implementation and testing of a digital pen and paper tool to support patients with heart failure and their health care providers in detecting early signs of deterioration and monitor adherence. *Int J Integr Care* 2011 Jul 21;11(7):e115. [doi: [10.5334/ijic.787](#)]
85. Maddocks H, Marshall JN, Stewart M, Terry AL, Cejic S, Hammond J, et al. Quality of congestive heart failure care: Assessing measurement of care using electronic medical records. *Can Fam Physician* 2010 Dec;56(12):e432-e437 [[FREE Full text](#)] [Medline: [21156884](#)]
86. Korb K, Hummers-Pradier E, Stich K, Chenot J, Scherer M. Implementation of recommendations for the diagnosis of heart failure. [corrected] [Article in German]. *Dtsch Med Wochenschr* 2010 Jan;135(4):120-124. [doi: [10.1055/s-0029-1244827](#)] [Medline: [20101554](#)]
87. Fonarow GC, Albert NM, Curtis AB, Stough WG, Gheorghiade M, Heywood JT, et al. Improving evidence-based care for heart failure in outpatient cardiology practices: Primary results of the Registry to Improve the Use of Evidence-Based Heart Failure Therapies in the Outpatient Setting (IMPROVE HF). *Circulation* 2010 Aug 10;122(6):585-596. [doi: [10.1161/CIRCULATIONAHA.109.934471](#)] [Medline: [20660805](#)]
88. Vercauteren S, Castadot M, Vanhalewyn M, Boulanger S, Robert A, Bouvy C, et al. BELGIUM-HF (Better Efficacy in Lowering events by General practitioner's Intervention Using remote Monitoring in Heart Failure): Concept and feasibility. In: Proceedings of the Belgian Society of Cardiology, 28th Annual Scientific Meeting. 2009 Feb Presented at: Belgian Society of Cardiology, 28th Annual Scientific Meeting; January 29-31, 2009; Brussels, Belgium p. 117. [doi: [10.2143/AC.64.1.2034372](#)]
89. Majeed A, Williams J, de Lusignan S, Chan T. Management of heart failure in primary care after implementation of the National Service Framework for Coronary Heart Disease: A cross-sectional study. *Public Health* 2005 Feb;119(2):105-111. [doi: [10.1016/j.puhe.2004.06.006](#)] [Medline: [15694957](#)]
90. Subramanian U, Fihn SD, Weinberger M, Plue L, Smith FE, Udris EM, et al. A controlled trial of including symptom data in computer-based care suggestions for managing patients with chronic heart failure. *Am J Med* 2004 Mar 15;116(6):375-384. [doi: [10.1016/j.amjmed.2003.11.021](#)] [Medline: [15006586](#)]
91. Roth A, Kajiloti I, Elkayam I, Sander J, Kehati M, Golovner M. Telecardiology for patients with chronic heart failure: The 'SHL' experience in Israel. *Int J Cardiol* 2004 Oct;97(1):49-55. [doi: [10.1016/j.ijcard.2003.07.030](#)] [Medline: [15336806](#)]
92. Gnani S, Gray J, Khunti K, Majeed A. Managing heart failure in primary care: First steps in implementing the National Service Framework. *J Public Health (Oxf)* 2004 Mar;26(1):42-47. [Medline: [15044573](#)]
93. Grypdonck L, Aertgeerts B, Luyten F, Wollersheim H, Bellemans J, Peers K, et al. Development of quality indicators for an integrated approach of knee osteoarthritis. *J Rheumatol* 2014 Jun;41(6):1155-1162. [doi: [10.3899/jrheum.130680](#)] [Medline: [24737907](#)]

94. Jansen MJ, Hendriks EJ, Oostendorp RA, Dekker J, De Bie RA. Quality indicators indicate good adherence to the clinical practice guideline on "Osteoarthritis of the hip and knee" and few prognostic factors influence outcome indicators: A prospective cohort study. *Eur J Phys Rehabil Med* 2010 Sep;46(3):337-345 [[FREE Full text](#)] [Medline: [20926999](#)]
95. MacLean CH, Saag KG, Solomon DH, Morton SC, Sampsel S, Klippel JH. Measuring quality in arthritis care: Methods for developing the Arthritis Foundation's quality indicator set. *Arthritis Rheum* 2004 Apr 15;51(2):193-202 [[FREE Full text](#)] [doi: [10.1002/art.20248](#)] [Medline: [15077259](#)]
96. Peat G, Lawton H, Hay E, Greig J, Thomas E, KNE-SCI Study Group. Development of the Knee Standardized Clinical Interview: A research tool for studying the primary care clinical epidemiology of knee problems in older adults. *Rheumatology (Oxford)* 2002 Oct;41(10):1101-1108. [Medline: [12364627](#)]
97. Glasziou P, Irwig L, Mant D. Monitoring in chronic disease: A rational approach. *BMJ* 2005 Mar 19;330(7492):644-648 [[FREE Full text](#)] [doi: [10.1136/bmj.330.7492.644](#)] [Medline: [15774996](#)]
98. van Steenkiste BC, Jacobs JE, Verheijen NM, Levelink JH, Bottema BJ. A Delphi technique as a method for selecting the content of an electronic patient record for asthma. *Int J Med Inform* 2002 Apr;65(1):7-16. [Medline: [11904244](#)]
99. Gensichen J, Peitz M, Torge M, Mosig-Frey J, Wendt-Hermainski H, Rosemann T, et al. The "Depression Monitoring list" (DeMoL) with integrated PHQ-D-Rationale and design of a tool for the case management for depression in primary care [Article in German]. *Z Arztl Fortbild Qualitatssich* 2006;100(5):375-382. [Medline: [16955623](#)]
100. Loughheed MD, Taite A, ten Hove J, Morra A, Van Dam A, Ducharme FM, et al. Pan-Canadian asthma and COPD standards for electronic health records: A Canadian Thoracic Society Expert Working Group Report. *Can J Respir Crit Care Sleep Med* 2018 Nov 05;2(4):244-250. [doi: [10.1080/24745332.2018.1517623](#)]
101. Chmiel C, Birnbaum B, Gensichen J, Rosemann T, Frei A. The diabetes traffic light scheme: Development of an instrument for the case management in patients with diabetes mellitus in primary care [Article in German]. *Praxis (Bern 1994)* 2011 Nov 30;100(24):1457-1473. [doi: [10.1024/1661-8157/a000751](#)] [Medline: [22124957](#)]
102. Roland M. Linking physicians' pay to the quality of care: A major experiment in the United Kingdom. *N Engl J Med* 2004 Sep 30;351(14):1448-1454. [doi: [10.1056/NEJMp041294](#)] [Medline: [15459308](#)]
103. Chmiel C, Senn O, Rosemann T, Del Prete V, Steurer-Stey C. CoCo trial: Color-coded blood pressure Control, a randomized controlled study. *Patient Prefer Adherence* 2014;8:1383-1392 [[FREE Full text](#)] [doi: [10.2147/PPA.S68213](#)] [Medline: [25346595](#)]
104. Schectman G, Barnas G, Laud P, Cantwell L, Horton M, Zarling EJ. Prolonging the return visit interval in primary care. *Am J Med* 2005 Apr;118(4):393-399. [doi: [10.1016/j.amjmed.2005.01.003](#)] [Medline: [15808137](#)]
105. Bakker K, Apelqvist J, Schaper NC, International Working Group on Diabetic Foot Editorial Board. Practical guidelines on the management and prevention of the diabetic foot 2011. *Diabetes Metab Res Rev* 2012 Feb;28 Suppl 1:225-231. [doi: [10.1002/dmrr.2253](#)] [Medline: [22271742](#)]
106. Clarson LE, Nicholl BI, Bishop A, Edwards JJ, Daniel R, Mallen CD. Monitoring osteoarthritis: A cross-sectional survey in general practice. *Clin Med Insights Arthritis Musculoskelet Disord* 2013;6:85-91 [[FREE Full text](#)] [doi: [10.4137/CMAMD.S12606](#)] [Medline: [24324351](#)]
107. Scalvini S, Martinelli G, Baratti D, Domenighini D, Benigno M, Paletta L, et al. Telecardiology: One-lead electrocardiogram monitoring and nurse triage in chronic heart failure. *J Telemed Telecare* 2005;11 Suppl 1:18-20. [doi: [10.1258/1357633054461750](#)] [Medline: [16035981](#)]
108. Morilla-Herrera JC, Garcia-Mayor S, Martín-Santos FJ, Kaknani Uttumchandani S, Leon Campos Á, Caro Bautista J, et al. A systematic review of the effectiveness and roles of advanced practice nursing in older people. *Int J Nurs Stud* 2016 Jan;53:290-307. [doi: [10.1016/j.ijnurstu.2015.10.010](#)] [Medline: [26542652](#)]
109. Agrinier N, Altieri C, Alla F, Jay N, Dobre D, Thilly N, et al. The ICALOR home-nurse led disease management programme in heart failure prevents hospital readmission: A French nationwide time-series comparison. In: *Proceedings of the European Society of Cardiology Congress 2012*. 2012 Presented at: European Society of Cardiology Congress 2012; August 25-29, 2012; Munich, Germany p. 159-160 URL: https://academic.oup.com/eurheartj/article-pdf/33/suppl_1/19/1188353/ehs281.pdf

Abbreviations

AWMF: Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften eV
CCM: chronic care model
EMR: electronic medical record
HbA_{1c}: hemoglobin A_{1c}
QOF: Quality and Outcomes Framework

Edited by G Eysenbach; submitted 06.05.18; peer-reviewed by R Bush, M Reading, W Zhang; comments to author 01.09.18; revised version received 20.03.19; accepted 07.04.19; published 17.05.19

Please cite as:

Falck L, Zoller M, Rosemann T, Martínez-González NA, Chmiel C

Toward Standardized Monitoring of Patients With Chronic Diseases in Primary Care Using Electronic Medical Records: Systematic Review

JMIR Med Inform 2019;7(2):e10879

URL: <http://medinform.jmir.org/2019/2/e10879/>

doi: [10.2196/10879](https://doi.org/10.2196/10879)

PMID:

©Leandra Falck, Marco Zoller, Thomas Rosemann, Nahara Anani Martínez-González, Corinne Chmiel. Originally published in JMIR Medical Informatics (<http://medinform.jmir.org>), 17.05.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Medical Informatics, is properly cited. The complete bibliographic information, a link to the original publication on <http://medinform.jmir.org/>, as well as this copyright and license information must be included.

Appendix 1

Search strategy for (OVID) MEDLINE*.

#	Searches	Results
1	(exp Medical Records/ or ((medical or patient) adj3 (record* or file)).ab,ti.) and (general practitioners/ or physicians, family/ or exp Primary Health Care/ or ((general or family or primary) adj3 (doctor* or physician* or practi*)).ab,ti.) and (exp Automatic Data Processing/ or indicator.ab,ti. or managment.ab,ti. or monitoring.ab,ti. or parameter.ab,ti. or ((clinical or laboratory) adj3 (data or assess* or finding* or observ* or monitor* or examin* or check* or control*)).ab,ti. or ((data or information) adj6 (set or extract* or analys* or retriev* or yield* or gather*)).ab,ti. or (data adj6 (standard* or normali* or form*)).ab,ti. or ("data from" adj6 (file or record)).ab,ti.)	2088
2	limit 1 to (yr="2000 -Current" and (english or german))	1593
3	exp Diabetes Mellitus, Type 2/	91561
4	2 and 3	74
5	(self monitoring or (monitoring adj3 glucose)).ab,ti.	9074
6	4 not 5	69
7	exp Hypertension/	215840
8	2 and 7	79
9	exp Heart Failure/	91040
10	2 and 9	22
11	exp heart failure/ or exp myocardial ischemia/	442513
12	2 and 11	60
13	12 not 10	38
14	exp Asthma/	108981
15	2 and 14	37
16	exp Arthritis/	209132
17	2 and 16	16

*Similar search strategies were applied in Embase and Cochrane.

Appendix 2

List of all included studies including monitoring indicators for the five chronic conditions.

First author	Year of publication	Country of origin	Guideline used in the publication	Country of origin of guideline	Reference number
Diabetes mellitus					
Suija	2015	Estonia	Organisation for Economic Co-operation and Development (OECD), American Diabetes Association (ADA), World Health Organization WHO, European Society of Cardiology (ESC), National Institute for Health and Care Excellence (NICE)	various	[10]
Shah	2015	United Kingdom	unknown		[11]
Devkota	2015	USA	American Diabetes Association (ADA)	USA	[12]
Barkhuysen	2014	Netherlands	Dutch College of General Practitioners	Netherlands	[13]
Szzech	2014	USA	Kidney Disease Outcomes Quality Initiative (KDOQI)	USA	[14]
van Melle	2014	Netherlands	Dutch College of General Practitioners	Netherlands	[15]
Djalali	2014	Switzerland	Quality and Outcomes Framework indicator (QOF)	United Kingdom	[16]
Goff	2014	USA	unknown		[17]
Vidal-Pardo	2013	Spain	"Plan de Saude" (Health plan of Galicia)	Spain	[18]
Sidorenkov	2013	Netherlands	Quality and Outcomes Framework indicator (QOF), Dutch College of General Practitioners	United Kingdom, Netherlands	[19]
Winkley	2013	United Kingdom	unknown		[20]

Gavran	2012	Bosnia/Herzegovina	The Committee for Practice Guidelines (CPG)	Europe	[21]
Knudsen	2012	Denmark	Danish National guidelines	Denmark	[22]
Mata-Cases	2012	Spain	RedGDPS (Spain)	Spain	[23]
Nouwens	2012	Netherlands	Dutch College of Family Physicians	Netherlands	[24]
Satman	2012	Turkey	Standard Diabetes Management Procedures of Turkey	Turkey	[25]
Marley	2012	Australia	unknown		[26]
Patapas	2012	Canada	unknown		[27]
Staff	2012	Australia	United Kingdom Prospective Diabetes Study	United Kingdom	[28]
Hill	2012	Ireland	indicators selected by authors		[29]
Alfadda	2011	Saudi Arabia	American Diabetes Association (ADA)	USA	[30]
Dickerson	2011	USA	Diabetes Recognition Program (DRP) by the National Committee for Quality Assurance (NCQA)	USA	[31]
Vidal Pardo	2011	Spain	"Plan de Saude" (Health plan of Galicia)	Spain	[32]
Weenink	2011	Netherlands	unknown		[33]
Gladstone	2011	USA	Canadian Practice Guidelines	Canada	[34]
Holbrook	2011	Canada	unspecific cardiovascular risk factors		[35]
O'Connor	2011	USA	unknown		[36]
Sundquist	2011	SWE	Swedish National Guidelines	Sweden	[37]
Reddy	2010	Australia	Quality and Outcomes Framework indicator (QOF)	United Kingdom	[38]
Samoutis	2010	Cyprus	American Diabetes Association (ADA), St. Vincent Declaration	USA, Europe	[39]

Shah	2010	Canada	Canadian Diabetes Association	Canada	[40]
Petrazzuoli	2010	Italy	Quality and Outcomes Framework indicator (QOF)	United Kingdom	[41]
Sperl-Hillen	2010	USA	American Diabetes Association (ADA)	USA	[42]
Pedersen	2009	Greenland	Danish National Indicator Project	Denmark	[43]
Holbrook	2009	Canada	Canadian Diabetes Association and American Diabetes Association (ADA)	Canada, USA	[44]
Moharram	2008	Saudi Arabia	Canadian Diabetes Association	Canada	[45]
Novo	2008	Bosnia/Herzegovina	Canadian Diabetes Association	Canada	[46]
Samuels	2008	USA	American Diabetes Association (ADA)	USA	[47]
Smith	2008	USA	American Diabetes Association (ADA), National Committee for quality assurance	USA	[48]
Voorham	2008	NL	National Guidelines of Netherlands	Netherlands	[49]
Wens	2007	Belgium	various guidelines	Belgium, Netherlands, Germany, United Kingdom, France	[50]
Nitiyanant	2007	Thailand	unknown		[51]
Herrin	2006	USA	National Diabetes Quality Improvement Alliance guidelines	USA	[52]
Wan	2006	Australia	Australian guidelines for diabetes management in general practice, National	Australia, United Kingdom	[53]

			Institute for Health and Care Excellence (NICE)		
Al Khaja	2005	Bahrain	The Seventh Report of the Joint National Committee (JNC-7)	USA	[54]
Cueto-Manzano	2005	Mexico	unknown		[55]
Lusignan	2005	United Kingdom	National Institute for Health and Care Excellence (NICE)	United Kingdom	[56]
Sequeira	2004	Bahrain	The Sixth Report of the Joint National Committee (JNC-6)/International Society of Hypertension (ISH)	USA, international	[57]
Wermeille	2004	Switzerland	unknown		[58]
Goudswaard	2003	Netherlands	Dutch College of General Practitioners	Netherlands	[59]
Campbell	2002	United Kingdom	Developed within study		[60]
Parchman	2002	USA	American Diabetes Association (ADA)	USA	[61]
Renders	2001	Netherlands	Dutch College of General Practitioners	Denmark	[62]
Linmans	2001	Netherlands	unknown		[63]
Asthma					
Minard	2014	Canada	Asthma Care Map	Canada	[64]
Lim	2012	USA	National Assessment of Educational Progress (NAEP)	USA	[65]
Lougheed	2012	Canada	Canadian Asthma Consensus Guidelines/Canadian Thoracic Society	Canada	[66]
Oei	2011	Australia	unknown		[67]
Nokela	2010	Sweden	unknown		[68]
Yawn	2008	USA	APGAR tool/ National Heart, Lung, and Blood Institute (NHLBI)	USA	[69]

Baddar	2006	Oman	Oman's national Manual for the Management of Asthma in Adults	Oman	[70]
Campbell	2002	United Kingdom	indicators selected by authors		[60]
Arterial hypertension					
Suija	2015	Estonia	Organisation for Economic Co-operation and Development (OECD), European Society of Cardiology (ESC), World Health Organization (WHO)/ International Society of Hypertension (ISH)	international, Europe	[10]
Hasselstrom	2014	Sweden	unknown		[71]
Tong	2012	Malaysia	Clinical Practice Guideline for Hypertension in Malaysia	Malaysia	[72]
Holbrook	2011	Canada	unknown		[35]
Samoutis	2010	Cyprus	Report of the Joint National Committee (JNC), the European guidelines on cardiovascular disease prevention and the European Society of Hypertension, European Society of Cardiology Guidelines.	USA, Europe	[39]
Pavlik	2009	USA	The Seventh Report of the Joint National Committee (JNC-7)	USA	[73]
Chan	2006	China	Eli Lilly National clinic audit center, Report of the Joint National Committee (JNC)	United Kingdom, USA	[74]
Asnani	2005	Jamaica	Ministry of Health	Jamaica	[75]
Rabinowitz	2005	Israel	American Heart Association	USA	[76]

			(AHA)/American College of Cardiology (ACC), National Cholesterol Education Program		
Mitchell	2005	United Kingdom	unknown		[77]
Alli	2005	Italy	unknown		[78]
Tierney	2004	USA	unknown		[79]
Lackland	2004	USA	The Seventh Report of the Joint National Committee (JNC-7)	USA	[80]
Frijling	2003	Netherlands	Dutch college of General Practitioners	Netherlands	[81]
Heart failure					
Amarasingham	2013	USA	unknown		[82]
Logeart	2013	France	European Society of Cardiology (ESC)	Europe	[83]
Weenink	2011	Netherlands	unknown		[33]
Lind	2011	Sweden	unknown		[84]
Korb	2010	Germany	Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin (DEGAM)-Leitlinie	Germany	[86]
Maddocks	2010	Canada	Heart failure management incentive (ministry of health and long-term care)	Canada	[85]
Fonarow	2010	USA	American College of Cardiology/American Heart Association	USA	[87]
Vercauteren	2009	Belgium	unknown		[88]
Majeed	2005	United kingdom	Quality and Outcomes Framework indicator (QOF)	United Kingdom	[89]
Roth	2004	Israel	unknown		[91]
Subramanian	2004	USA	Kansas City Cardiomyopathy Questionnaire	USA	[90]

Gnani	2004	United Kingdom	Quality and Outcomes Framework indicator (QOF)	United Kingdom	[92]
Osteoarthritis					
Grypdonck	2014	Belgium	indicators selected by authors		[93]
Jansen	2010	Netherlands	Osteoarthritis of the hip and knee for physical therapist	Netherlands	[94]
MacLean	2004	USA	indicators selected by authors		[95]
Peat	2002	United Kingdom	indicators selected by authors		[96]

Appendix 3

Asthma indicators mentioned in guidelines and studies. The indicators are sorted first by guidelines and then by studies.

indicators for asthma	appeared in guidelines	appeared in studies
Fev ₁ /FVC Ratio	6 (a-f)	-
lung function	6 (a-f)	3 [64, 66, 69]
Daytime asthma symptoms	6 (a-f)	4 [60, 64, 66, 69]
Any night waking due to asthma	5 (a, c-f)	5 [60, 64, 66, 68, 69]
Any activity limitation due to asthma	5 (a, c-f)	4 [60, 64, 66, 69]
FEV ₁	5 (a-c, e, f)	1 [68]
PEF (self-monitoring)	5 (a-c, e, f)	4 [60, 64, 68, 70]
wheezing	4 (b-e)	2 [64, 69]
cough	4 (b-e)	2 [64, 69]
Reliever needed for symptoms more than twice per week	4 (a, c, d, f)	2 [64, 68]
smoking habit	4 (a-d)	3 [60, 65, 68]
auscultation	4 (a, b, d, e)	-
bronchial provocation test	4 (a, b, d, e)	-
variation in lung function	4 (a, d-f)	-
check inhaler technique	3 (a-c)	4 [60, 64, 69, 70]
trigger of symptoms	3 (a, c, d)	3 [64, 69, 70]
hospital stays or emergency department visits since last visit	1 (c)	3 [64, 66, 70]
medication history	2 (a,c)	3 [64, 69, 70]
smoking cessation advice	-	3 [60, 64, 66]
check adherence	2 (a,c)	3 [64, 68, 70]
asthma action plan	-	3 [64, 66, 69]
indicators appeared in less than 4 guidelines	162	

indicators appeared in less than 3 studies		38
--	--	-----------

Letters a-f refers to the guidelines listed in Appendix 8; FEV₁/FVC Ratio: Tiffeneau-Pinelli index.

FEV₁: Forced expiratory volume at one second. PEF: peak expiratory flow.

Appendix 4

Arterial hypertension indicators mentioned in guidelines and studies. The indicators are sorted first by guidelines and then by studies.

indicators for arterial hypertension	appeared in guidelines	appeared in studies
smoking habit	8 (a-h)	11 [35, 71-79, 81]
blood pressure	8 (a-h)	9 [10, 35, 39, 71-73, 75, 77-79]
choose correct cuff	8 (a-h)	1 [75]
standing blood pressure	8 (a-h)	-
electrocardiogramm	8 (a-h)	2 [72, 75]
fasting blood glucose and or HbA1c	8 (a-h)	6 [39, 71, 72, 75, 79, 80]
serum totalcholesterine	8 (a-h)	8 [10, 39, 71, 72, 74, 75, 79, 80]
high density lipoprotein	8 (a-h)	5 [39, 71, 72, 76, 80]
creatinine	8 (a-h)	4 [71, 75, 79, 80]
potassium	8 (a-h)	3 [75, 79, 80]
age	7 (a-f, h)	2 [76, 78]
history of chronic kidney disease	7 (a, c-h)	2 [78, 79]
recreational drug use	7 (a, c-h)	-
current medication	7 (a, c-h)	3 [73, 75-77]
family history	7 (a-d, g, h)	3 [72, 74, 76]
family history for coronary heart disease	7 (a-d, g, h)	3 [72, 74, 76]
BMI/weight	7 (a-f, h)	8 [35, 71, 72, 74-76, 78, 79]
resting comfortably for several minutes prior to blood pressure measurement	7 (a, b, d-h)	-
measure blood pressure on both arms	7 (a-e, g,h)	-
Home blood pressure monitoring or ambulatory blood pressure monitoring if suspected white coat hypertension	7 (a-g)	-
eye examination	7 (a-d, e-h)	3 [72, 74, 75]
echocardiography if needed	7 (a-c, e-h)	-

low density lipoprotein	7 (a, b, d-h)	6 [35, 39, 71, 72, 76, 80]
triglycerides	7 (a, b, d-h)	6 [39, 71, 72, 75, 76, 80]
Assessment of microalbuminuria	7 (a, b, d-h)	4 [71, 72, 79, 80]
sedentary lifestyle	6 (a, d-h)	5 [35, 74-76, 78]
diabetes mellitus in history	6 (a, c, d, f-h)	6 [72, 74, 76-79]
alcohol	3 (e, g, h)	4 [74, 75, 78, 79]
history of coronary heart disease/cardiovascular disease/stroke	5 (a, d-g)	2 [77-79]
indicators appeared in less than 7 guidelines	307	
indicators appeared in less than 3 studies		58

Letters a-h refer to the guidelines listed in Appendix 9; BMI: body mass index.

Appendix 5

Chronic heart failure indicators most frequently mentioned in guidelines and studies. The indicators are sorted first by guidelines and then by studies.

indicators for chronic heart failure	appeared in guidelines	appeared in studies
sodium	6 (a-f)	4 [33, 85, 87, 89]
potassium	6 (a-f)	3 [33, 85, 89]
creatinine	6 (a-f)	5 [33, 82, 83, 85, 87]
daily weight measurement	5 (a, b, d-f)	3 [84, 88, 91]
changes in weight	5 (a, b, d-f)	
depression	5 (a, c-f)	-
address palliative or hospice care	5 (a, c, d-e)	-
NYHA-classification	5 (a, c-f)	2 [87, 90]
pulse rate	5 (a-c, e-f)	3 [83, 85, 87]
exercise capacity	5 (a, b, d-f)	2 [86, 90]
erectile dysfunction/sexual activity	4 (a, d, e, f)	-
measure blood pressure	4 (a, b, d, e)	5 [33, 85, 87, 89, 92]
measure daily blood pressure		3 [84, 88, 91]
electrocardiography	4 (a, d-f)	4 [85, 86, 89, 92]
echocardiogram with doppler	4 (a, d-f)	6 [85-89, 92]
NT-proBNP	3 (d-f)	5 [82, 83, 86-88]
chest radiography if necessary	3 (d-f)	4 [85, 86, 89, 92]
BMI	2 (e, f)	3 [33, 85, 89]
history/signs of diabetes mellitus	3 (a, d, f)	3 [82, 87, 89]
number of indicators that appeared in less than 4 guidelines	213	
number of indicators that appeared in less than 3 studies		47

Letters a-f refer to the guidelines listed in Appendix 10; NT-proBNP: N-terminal prohormone Brain natriuretic peptide. BMI: body mass index.

Appendix 6

Osteoarthritis indicators most frequently mentioned in guidelines and studies. The indicators are sorted first by guidelines and then by studies.

indicators for osteoarthritis	appeared in guidelines	appeared in studies
tenderness	4 (a-d)	
significant loss of range of movement	4 (a-d)	
assessment of functional status		2 [94, 95]
mood	3 (a-c)	
health believes	3 (a-c)	
expectations, concerns, ideas	3 (a-c)	
stiffness	3 (a, c, d)	
stress pain	3 (a, c, d)	
activity of daily living	3 (a, b, d)	
Hobbies	3 (a-c)	
history of trauma	3 (a, c, d)	
swelling	3 (a, c, d)	
redness	3 (a-c)	
deformity	3 (a, b, d)	
advise on losing and maintaining weight		2 [93, 95]
assessment of pain		2 [94, 95]
number of indicators that appeared in less than 3 guidelines	148	
number of indicators that appeared in only one study		38

Letters a-f refer to the guidelines listed in Appendix 11.

Appendix 7

Guidelines screened for indicators for diabetes mellitus type2.

Diabetes mellitus	Year (last update)	editor/publisher	country	
Standards of Medical Care in Diabetes	2014	American Diabetes Association (ADA)	USA	a
Nationale VersorgungsLeitlinien	2013, 2015	Bundesärztekammer (BÄK), Kassenärztliche Bundesvereinigung (KBV), Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF) (AWMF Institute for Medical Knowledge Management)	Germany	b
National Institute for Health and Care Excellence (NICE)	2014	Royal College of Physicians	England	c
Global Guideline for Type 2 Diabetes	2012	International Diabetes Federation	International	d
General practice management of type 2 diabetes	2014	The Royal Australian College of General Practitioners	Australa	e
Clinical Practice Guidelines	2013	Canadian Diabetes Association	Canada	f
MediX-Guideline zu Diabetes mellitus	2013	MediX	Schweiz	g

Appendix 8

Guidelines screened for indicators for asthma.

Asthma	Year (last update)	editor/publisher	country	
Global strategy for Asthma Management and Prevention	2015	Global Initiative for Asthma (GINA)	international	a
Asthma: diagnosis and monitoring of asthma in adults, children and young people	Draft for Consultation 2015	National Institute for Health and Care Excellence (NICE)	England	b
Guidelines for the Diagnosis and Management of Asthma	2007	National Heart, Lung, and Blood Institute (NHBLI)	USA	c
Australian Asthma Handbook (Quick Reference Guide)	2014	The Royal Australian College of General Practitioners	Australia	d
Leitlinien zu Diagnostik und Therapie von Patienten mit Asthma	2006	Deutsche Atemwegsliga, Deutsche Gesellschaft für Pneumologie und Beatmungsmedizin e.V.	Germany	e
MediX-Guideline zu Asthma bronchiale	2011	MediX	Switzerland	f

Appendix 9

Guidelines screened for indicators for arterial hypertension.

Arterial hypertension	Year (last update)	editor/publisher	country	
Leitlinien für das Management der arteriellen Hypertonie (Guidelines for management of arterial hypertension)	2013	Deutsche Gesellschaft für Kardiologie-, Herz- und Kreislaufforschung (DGK) (German Cardiac Society) and Deutsche Hochdruckliga e.V. DHL® Deutsche Gesellschaft für Hypertonie und Prävention	Germany	a
Practice Guidelines for the Management of Arterial Hypertension	2007	The European Society of Cardiology (ESC) and European Society of Hypertension (ESH)	Europe	b
Clinical management of primary hypertension in adults	2013	National Institute for Health and Clinical Excellence (NICE)	England	c
The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure	2003	National Heart, Lung, and Blood Institute (NHLBI)	USA	d
The 2015 Canadian Hypertension Education Program Recommendations	2015	Hypertension Canada	Canada	e
Clinical Practice Guidelines for the Management of Hypertension in the Community	2014	The American Society of Hypertension and the International Society of Hypertension (ISH)	USA, international	f
Guide to management of hypertension 2008 (Assessing and managing raised blood pressure in adults)	2010	National Heart Foundation of Australia	Australia	g

Arterielle Hypertonie Empfehlungen für Ärzte	2015	Swiss Society of Hypertension (Schweizerische Hypertonie Gesellschaft)	Switzerland	h
--	------	--	-------------	---

Appendix 10

Guidelines screened for indicators for chronic heart failure.

Chronic heart failure	Year (last update)	editor/publisher	country	
Nationale VersorgungsLeitlinie	2011	Bundesärztekammer (BÄK), Kassenärztliche Bundesvereinigung (KBV), Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF) (AWMF Institute for Medical Knowledge Management)	Germany	a
Anforderungen an ein Modul „Chronische Herzinsuffizienz“ für strukturierte Behandlungsprogramme für Koronare Herzkrankheit (KHK)	--	Empfehlungen des Gemeinsamen Bundesausschusses gemäss §137f Abs. 2 SGB V für die Rechtsverordnung nach § 266 Abs. 7 SGB V	Germany	b
Chronic Heart Failure; National clinical guideline for diagnosis and management in primary and secondary care	2010	National Institute for Health and Clinical Excellence (NICE)	England	c
ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012	2012	The European Society of Cardiology (ESC)	Europe	d
Guideline for the Management of Heart Failure	2013	The American College of Cardiology Foundation and the American Heart Association	USA	e
Guidelines for the prevention, detection and management of chronic heart failure in Australia	2011	National Heart Foundation of Australia	Australia	f

Appendix 11

Guidelines screened for indicators for osteoarthritis.

Osteoarthritis	Year (last update)	editor/publisher	country	
Osteoarthritis; Care and management in adults	2014	National Institute for Health and Clinical Excellence (NICE)	England	a
EULAR recommendations for the non-pharmacological core management of hip and knee osteoarthritis	2013	The European League against Rheumatism	Europe	b
Guideline for the non-surgical management of hip and knee osteoarthritis	2009	The Royal Australian College of General Practitioners	Australia	c
MediX-Guideline zu Arthrose	2013	MediX	Schweiz	d